

ach in an hour, and rarely continuing longer than four. He found it to excite little or no nausea previous to its operation, and though it made a powerful impression on the system, it never produced any disagreeable or unusual symptoms.

Dr. Hayward also made trial of the powder of the leaves, which he found to possess the same properties with that of the root, but to be less effectual and less certain in its operation. He also prepared a tincture, decoction, and wine of the root; but all these were inferior to the medicine in substance, being less certain in their effect, and sometimes giving rise to troublesome symptoms.

Dr. Shultz of Pennsylvania, author of an inaugural dissertation on the *Phytolacca decandra*, gave the expressed juice of the leaves, berries, and roots, in considerable quantity to animals. It operated by emesis and catharsis, attended with drowsiness. The juice of the root was most active. He also gave to a dog two ounces of the spiritous liquor distilled from the berries. It occasioned nausea and drowsiness, with slight spasmodic motions, but no vomiting.

In the same dissertation, Dr. Shultz refers to several instances of persons who had incautiously eaten large quantities of the root through mistake. Its effects were violent vomiting and purging,

prostration of strength, and in some instances convulsions.

The *Phytolacca* has had some reputation in the treatment of rheumatism. Dr. Griffiths, formerly a professor in the University of Pennsylvania, found it of great use in Syphilitic rheumatism. Dr. Hayward however states, that he derived no advantage from its employment in rheumatic affections.

The young shoots of this vegetable are destitute of medicinal qualities, and are eaten in the spring in some parts of the United States, as substitutes for asparagus. At this time the *succus proprius* or returning juice of the plant is not yet formed by exposure of the sap to the atmospheric air, in the leaves. The ripe berries are less noxious than the green, and are devoured by several species of birds. In Portugal and in France they were formerly employed to improve the colour of red wines, until the interference of government became necessary to put a stop to the practice.

The external application of *Phytolacca* has been found useful in a variety of cases, by its action as a local stimulant. The ointment and extract have commonly been employed for this purpose. These preparations usually excite a sense of